


Recommended sites for childhood vaccinations in NSW

JULY 2020
SCHEDULE

6 WEEKS AND 4 MONTHS




Bexsero® (IM)
Aboriginal† Children only

Rotarix® (ORAL)
Check age limit for Dose 1 and Dose 2

Infanrix Hexa® (IM)

Prevenar 13® (IM)

6 MONTHS




Annual Influenza funded for:

- All children ≥6 months to less than 5 years of age.
- All Aboriginal† people ≥6 months
- All people ≥6 months with at risk medical.


- At risk children ≥6 months require an additional dose of Prevenar 13® (pneumococcal) - refer to AIH* online.
- Aboriginal† children ≥6 months with certain medical conditions may require an additional dose of Meningococcal B - refer to AIH* online.

12 MONTHS




- Additional dose of Hepatitis B vaccine recommended at 12 months for Low birth weight (<2000gms) and/or infants born at <32 weeks gestation - refer to AIH* online.

18 MONTHS



- Additional dose of Hepatitis B vaccine recommended at 12 months for Low birth weight (<2000gms) and/or infants born at <32 weeks gestation - refer to AIH* online.

4 YEARS



- At risk children require a dose of Pneumovax 23® (pneumococcal) at 2-12 months after Prevenar 13® or at age 4 years whichever is later.

Before Vaccination:

- Adrenaline Kit readily available.
- Confirm vaccine cold chain maintained.
- Pre-vaccination checklist attended.
- Identify all medically at risk children and Aboriginal† children – requiring additional vaccines as per AIH* online.
- Check child’s immunisation history on AIR^.
- Informed consent obtained and documented.
- Check if the vaccine needs to be reconstituted.
- Ensure correct vaccine for age and give all schedule vaccines in one visit if possible.

Post Vaccination:


- Child remains in clinic for 15 minutes.
- Provide parent/carer with Handout - *What to Expect & What to Do?*
- Add recall reminder for next visit.


Bexsero®


- Children <2 years of age are recommended to receive prophylactic paracetamol 30 minutes before or as soon as practical after Bexsero® vaccination.
- Advise parent/carer to follow up with 2 more doses of paracetamol given 6 hours apart, regardless of whether the child has a fever.
- Children <2 years of age can receive Bexsero® with other scheduled vaccines.
- Children <2 years of age can receive Bexsero® if required separately from other infant vaccines, with a minimum interval of 3 days.

At Risk Groups

- MenB, MenACWY, pneumococcal and/or HIB vaccines are funded for all people with:
 - Asplenia
 - Hyposplenia
 - Complement deficiency
 - Treatment with Eculizumab
- Refer to AIH* online.

INTRAMUSCULAR (IM) 90° to skin plane		
	Infant or Child	22–25 gauge, 25 mm long
	Preterm infant (<37 weeks gestation) up to 2 months of age, and/ or very small infant	23–25 gauge, 16 mm long

SUBCUTANEOUS (SC) 45° to skin plane		
	Infant or Child	25–27 gauge, 16 mm long



ADMINISTERING 3-4 VACCINES AT ONE VISIT

- Ensure 2.5cm between injections in the same muscle.
- If the child is receiving 3 or 4 vaccines at 12 months or greater and deltoid muscles mass is inadequate, give the less reactogenic vaccines (e.g. MMR and/or MenACWY) into the anterolateral thigh.

Aboriginal†: is inclusive of Aboriginal and Torres Strait Islander people
AIH* online: Australian Immunisation Handbook online
AIR^: Australian Immunisation Register

Murrumbidgee PHN and Murrumbidgee LHD developed this resource in April 2019 based on resources from Tasmania PHN, Hunter New England LHD and Central & Eastern Sydney PHN and updated July 2020. Document endorsed by Western NSW Local Health District and Far West Local Health District for distribution by Western NSW Primary Health Network.

phn
WESTERN NSW
An Australian Government Initiative

NSW
GOVERNMENT